

# End of life, preferred place of death (Retired)

## Alternate Name\*

End of life, preferred place of death

## INDICATOR DESCRIPTION

---

### Description\*

This indicator measures the percentage of palliative/end of life patients who died in their preferred place of death.

### Indicator Status\*

Retired

### HQO Reporting tool/product

Quality Improvement Plans (QIPs)

### Dimension\*

Patient-centred

### Type\*

Outcome

## DEFINITION AND SOURCE INFORMATION

---

### Unit of Measurement\*

Percentage

### Calculation Methods\*

The percentage is calculated as: numerator divided by denominator times 100.

### Numerator including inclusion/exclusion\*

Number of palliative / end of life patients who died in their preferred place of death. Patient must be discharged with a disposition of death in the period selected

*Inclusions:*

- In home referrals only
- > 18 years of age at the time
- CCM population of “complex palliative” or “chronic palliative” or end of life (SRC95) at time of discharge
- Discharge dispositions:
  - Died at home (preferred place of death)
  - Died in hospital (preferred place of death)
  - Died in residential hospice (preferred place of death)
  - Died in Long-Term Care home (preferred place of death)
  - Died in Retirement Home (preferred place of death)
  - Died in other location (preferred place of death)

*Exclusions:*

- Other CCM or SRC codes
- Children &le;18 years old at the time of discharge
- Referrals other than in home (school, placement)
- Other death discharge dispositions (not preferred place of death)
- Patients with blank or invalid health card numbers

**Denominator including inclusion/exclusion\***

Number of palliative/ end of life patients whose discharge disposition is death in the reporting period.

*Inclusions:*

- In home referrals only

- > 18 years of age at the time
- CCM population of “complex palliative” or “chronic palliative” or end of life (SRC95) at time of discharge
- Any discharge where Discharge disposition flag “has died” = 1
- Died (while under LHIN Home and community care services)
- Died in Hospital (< 14 days)
- Died in Hospital
- Died at Home
- Died at Home (preferred place of death)
- Died in Hospital (preferred place of death)
- Died in Residential Hospice (preferred place of death)
- Died in LTCH (preferred place of death)
- Died in Retirement Home
- Died in Retirement Home (preferred place of death)
- Died in Other Location
- Died in Other Location (preferred place of death)
- Died in Residential Hospice
- Unplanned Death

*Exclusions:*

- Other CCM or SRC codes
- Children &le;18 years old at the time of discharge
- Referrals other than in home (school, placement)
- Discharges other than death
- Patients with blank or invalid health card numbers

### Data Source

Client Health and Related Information System (CHRIS)

### Data provided to HQO by

Health Shared Services Ontario (HSSO)

## OTHER RELEVANT INFORMATION

---

### Comments Detailed

This is a QIP priority indicator for 2018/19. QIP current performance reporting period: October 2016 - September 2017  
To access your organization's data for the reporting period, refer to Health Quality Ontario's QIP Navigator. Data will be available in February 2018. Alternatively, to access your organization's data for this indicator, refer to CCAC Reporting Portal. This indicator was retired in the 2019/20 QIP.

## TAGS

---

### TAGS\*

Home Care  
Outcome  
End-of-life / Palliative  
Patient-centred  
Client Health and Related Information System (CHRIS)

## PUBLISH

---

### PUBLISH DATETIME\*

04/03/2019 10:11:00