

# End of life, preferred place of death (Retired)

## Alternate Name\*

End of life, preferred place of death

## INDICATOR DESCRIPTION

---

### Description\*

This indicator measures the percentage of palliative/end of life patients who died in their preferred place of death.

### HQO Reporting tool/product

Quality Improvement Plans (QIPs)

### Dimension\*

Patient-centred

### Type\*

Outcome

## DEFINITION AND SOURCE INFORMATION

---

### Unit of Measurement\*

Percentage

### Calculation Methods\*

The percentage is calculated as: numerator divided by denominator times 100.

### Numerator (short description i.e. not inclusions/exclusions)\*

Number of palliative / end of life patients who died in their preferred place of death. Patient must be discharged with a disposition of death in the period selected.

### Denominator (short description i.e. not inclusions/exclusions)\*

Number of palliative/ end of life patients whose discharge disposition is death in the reporting period.

**Adjustment (risk, age/sex standardization)- generalized**

None

**Data Source**

Client Health and Related Information System (CHRIS)

**Data provided to HQO by**

Health Shared Services Ontario (HSSO)

---

**OTHER RELEVANT INFORMATION**

**Comments Summary**

This is a QIP priority indicator for 2018/19. QIP current performance reporting period: October 2016 - September 2017  
To access your organization's data for the reporting period, refer to Health Quality Ontario's QIP Navigator. Data will be available in February 2018. Alternatively, to access your organization's data for this indicator, refer to CCAC Reporting Portal. This indicator was retired in the 2019/20 QIP.

---

**TAGS**

**TAGS\***

Home Care  
Outcome  
End-of-life / Palliative  
Patient-centred  
Client Health and Related Information System (CHRIS)

---

**PUBLISH**

**PUBLISH DATETIME\***

04/03/2019 10:11:00